

Melioidose : Références

References

[1] Abbink FC, Orendi JM, de Beaufort AJ.

Mother-to-child transmission of *Burkholderia pseudomallei*.

N Engl J Med. 2001 ;344:1171-1172.

[2] Allen RK.

La mélioidose : un diagnostic à évoquer devant une pneumopathie au retour d'Océanie.

Rev Mal Respir. 2002 ;19:807-808.

[3] Appassakij H, Silpapojakul KR, Wansit R, Pornpatkul M.

Diagnostic value of the indirect hemagglutination test for melioidosis in an endemic area.

Am J Trop Med Hyg. 1990 ;42:248-253.

[4] Ashdown LR.

An improved screening technique for isolation of *Pseudomonas pseudomallei* from clinical specimens.

Pathology. 1979 Apr ;11(2):293-7.

[5] Brett PJ, Deshazer D, Woods DE.

Characterization of *Burkholderia pseudomallei* and *Burkholderia pseudomallei*-like strains.

Epidemiol Infect. 1997 ;118:137-148.

[6] Brett PJ, Woods DE.

Pathogenesis of and immunity to melioidosis.

Acta Trop. 2000 ;74:201-210.

[7] Buchet R, Tredaniel J.

La mélioidose pulmonaire.

Ann Radiol Paris. 1985 ;28:387-390.

[8] Campus de microbiologie médicale.

Fiches de diagnostic.

<http://www.uvp5.univ-paris5.fr/MICROBES/diagnostic/pseudomallei/pseudomallei.asp>

(consulté le 09.01.2003).

[9] Chadwick DR, Ang B, Sitoh YY, Lee CC.

Cerebral melioidosis in Singapore : a review of five cases.

Trans R Soc Trop Med Hyg. 2002 ;96:72-76.

[10] Chaowagul W.

Recent advance in treatment of severe melioidosis.

Acta Trop. 2000,74 : 133-137.

[11] Chaowagul W, White NJ, Dance DA, Wattanagoon Y, Naigowit P, Davis TM, et al.

Melioidosis : a major cause of community-acquired septicemia in northeastern Thailand.

J Infect Dis. 1989 ;159:890-899.

[12] Charuchaimontri C, Suputtamongkol Y, Nilakul C, Chaowagul W, Chetchotisakd P, Lertpatanasuwun N, et al.

Antilipopolysaccharide II : an antibody protective against fatal melioidosis.

Clin Infect Dis. 1999 ;29(4) :813-8.

[13] Cherian T, Raghupathy P, John TJ.

Plague in India.

The Lancet. 1995 ;345:258-259.

[14] Chetchotisakd P, Porramatikul S, Mootsikapun P, Anunnatsiri S, Thinkhamrop B.

Randomized, double-blind, controlled study of cefoperazone-sulbactam plus cotrimoxazole versus ceftazidime plus cotrimoxazole for the treatment of severe melioidosis.

Clin Infect Dis. 2001 ;33:29-34.

[15] Choy JL, Mayo M, Janmaat A, Currie BJ.

Animal melioidosis in Australia.

Acta Trop. 2000 ;74:153-158.

[16] Currie BJ, Fisher DA, Anstey NM, Jacups SP.

Melioidosis : acute and chronic disease, relapse and re-activation.

Trans R Soc Trop Med Hyg. 2000 ;94:301-304.

[17] Currie BJ, Fisher DA, Howard DM, Burrow JN.

Neurological melioidosis.

Acta Trop. 2000 ;74:145-151.

[18] Currie BJ, Fisher DA, Howard DM, Burrow JN, Lo D, Selva-Nayagam S et al.

Endemic melioidosis in tropical northern Australia : a 10-year prospective study and review of the literature.

Clin Infect Dis. 2000 ;31:981-986.

[19] Currie BJ, Fisher DA, Howard DM, Burrow JN, Selvanayagam S, Snelling PL et al.

The epidemiology of melioidosis in Australia and Papua New Guinea.

Acta Trop. 2000 ;74:121-127.

[20] Dance DA.

Ecology of *Burkholderia pseudomallei* and the interactions between environmental *Burkholderia* spp. and human-animal hosts.

Acta Trop. 2000 ;74:159-168.

[21] Dance Da.

Melioidosis as an emerging global problem.

Acta Trop. 2000 ;74:115-119.

[22] Dance DA.

Melioidosis : the tip of the iceberg ?

Clin Microbiol Rev. 1991 ;4:52-60.

[23] Dance DA, Sanders D, Pitt TL, Speller DC.

Burkholderia pseudomallei and Indian Plague-like illness.

The Lancet. 1995 ;346:904-905.

[24] Dance DA, Smith MD, Aucken HM, Pitt TL.

Imported melioidosis in England and Wales.

Lancet. 1999 ;353:208.

[25] Dance DA, Wuthiekanun V, Chaowagul W, White NJ.

The antimicrobial susceptibility of *Pseudomonas pseudomallei*.

Emergence of resistance in vitro and during treatment.

J Antimicrob Chemother. 1989 ;24:295-309.

[26] Dance DA, Wuthiekanun V, Naigowit P, White NJ.

Identification of *Pseudomonas pseudomallei* in clinical practice : use of simple screening tests and API 20NE.

J Clin Pathol. 1989 ;42:645-648.

[27] Desakorn V, Smith MD, Wuthiekanun V, Dance DA, Auckan H, Suntharasamai P et al.

Detection of *Pseudomonas pseudomallei* antigen in urine for the diagnosis of melioidosis.

Am J Trop Med Hyg. 1994 ;51:627-633.

[28] Dharakul T, Songsivilai S.

The many facets of melioidosis.

Trends Microbiol. 1999 ;7:138-140.

[29] Dhiensiri T, Eua-Ananta Y.

Visceral abscess in melioidosis.

J Med Assoc Thai. 1995 ;78:225-231.

[30] Dhiensiri T, Puapairoj S, Susaengrat W.

Pulmonary melioidosis : clinical-radiologic correlation in 183 cases in northeastern Thailand.

Radiology. 1988 ;166:711-715.

[31] Dodin A, Galimand M.

Mélioïdose.

Encycl Med Chir, Maladies Infectieuses, Paris, 8036-C-10,1985.

[32] Dodin A, Galimand M.

Naissance, vie et recul d'une maladie infectieuse. La mélioïdose en pays tempéré.

Arch Inst Pasteur Tunis. 1986 ;63:69-73.

[33] Dodin A, Galimand M, Chove MA, Sanson R.

Le bacille de Whitmore. Germe d'actualité.

Med Mal Infect. 1976 ;6-10b:395-398.

[34] Dorman SE, Gill VJ, Gallin JI, Holland SM.

Burkholderia pseudomallei infection in a Puerto Rican patient with chronic granulomatous disease : case report and review of occurrences in the Americas.

Clin Infect Dis. 1998 ;26:889-894.

[35] Edmond KM, Bauert P, Currie BJ.

Paediatric melioidosis in the Northern Territory of Australia : an expanding clinical spectrum.

J Paediatr Child Health. 2001 ;37:337-341.

[36] Euzeby JP.

Burkholderia pseudomallei, Burkholderia thailandensis. Internet : Dictionnaire de bactériologie vétérinaire.

<http://www.bacterio.cict.fr/bacdico/garde.html>

(consulté le 09.01.2002).

[37] Galimand M.

Le bacille de Whitmore : caractérisation de Pseudomonas pseudomallei et physiopathologie de la mélioïdose.

Thèse de doctorat es Sciences, Université Paris 7, 1986 : 210p.

[38] Galimand M, Dodin A.

Le point sur la mélioïdose dans le monde.

Bull Soc Path Ex. 1982 ;75:375-383.

[39] Gamerman H, Mollaret HH, Dodin A, Delecloy JM, Romeo R.

Une maladie infectieuse qu'il faut savoir, désormais, évoquer en France : la mélioïdose.

Med Mal Infect. 1977 ;7:395-399.

[40] Gilligan PH.

Therapeutic challenges posed by bacterial bioterrorism threats.

Curr Opin Microbiol. 2002 ;5:489-495.

[41] Haase A, Janzen J, Barret S, Currie B.

Toxin production by Burkholderia pseudomallei strains and correlation with severity of melioidosis.

J Med Microbiol. 1997 ;46:557-53.

[42] Handa R, Bhatia S, Wali JP.

Melioidosis : a rare but not forgotten cause of fever of unknown origin.

Br J Clin Pract. 1996 ;50:116-117.

[43] Harle JR, Disdier P, Dodin A, Lebreuil G, Ottomani A, Weiller PJ.

Mélioïdose : forme aiguë avec des abcès spléniques.

Presse Med. 1990 ;19:1154.

[44] Heyse AM, Dierick J, Vanhouteghem H, Ameye F, Baert D, Burvenich P et al.

A case of imported melioidosis presenting as prostatitis.

Infection. 2003 ;31:60-62.

[45] Holland DJ, Wesley A, Drinkovic D, Currie BJ.

Cystic fibrosis and Burkholderia pseudomallei infection : an emerging problem ?

Clin Infect Dis. 2002 ;35(12) :e138-40.

[46] Hoque SN, Minassian M, Clipstone S, Lloyd-Owen SJ, Sheridan E, Lessing MP.

Melioidosis presenting as septic arthritis in Bengali men in east London. Rheumatology (Oxford). 1999 ;38:1029-1031.

[47] Hovette P, Aubron C.

Mélioïdose.

Encycl Med Chir, Maladies infectieuses. Paris : Éditions Scientifiques et Médicales Elsevier SAS,8-036-c-10,2001,8p.

[48] Hsueh PR, Teng LJ, Lee LN, Yu CJ, Yang PC, Ho SW et al.

Melioidosis : an emerging infection in Taiwan ?

Emerg Infect Dis. 2001 ;7:428-433.

[49] Hueck CJ.

Type III protein secretion systems in bacterial pathogens of animals and plants.

Microbiol Mol Biol Rev. 1998 Jun ;62:379-433.

[50] Husson MO, Hamze M, Verhille S et al.

Pseudomonas et *Burkholderia* in. Freney J, Renaud F, Hansen W et al. Précis de bactériologie clinique. Paris : Eska, 2000:1259-1285.

[51] IP M, Osterberg LG, Chau PY, Raffin TA.

Pulmonary melioidosis.

Chest.1995 ;108:1420-1424.

[52] Jaussaud R, Brasme L, Beguinot I, Vernet V, Rémy G, Deville J.

Une étiologie de fièvre au retour d'Asie du Sud-Est à ne pas méconnaître : la mélioiïdose.

Rev med Interne. 1997 ;18:201s.

[53] Jenney AW, Lum G, Fisher DA, Currie BJ.

Antibiotic susceptibility of *Burkholderia pseudomallei* from tropical northern Australia and implications for therapy of melioidosis.

Int J Antimicrob Agents. 2001 ;17:109-13.

[54] John TJ, Jesudason MV, Lalitha MK, Ganesh A, Mohandas V, Cherian T et al.

Melioidosis In India : the tip of the iceberg ?

Indian J Med Res. 1996 ;103:62-65.

[55] Leelarasamee A.

Melioidosis in Southeast Asia.

Acta Trop. 2000 ;74:129-132.

[56] Leelarasamee A, Bovornkitti S.

Melioidosis : review and update.

Rev Infect Dis. 1989 ;11:413-425.

[57] Lumbiganon P, Viengnondha S.

Clinical manifestations of melioidosis in children.

Pediatr Infect Dis J. 1995 ;14:136-140.

[58] McCormick JB, Sexton DJ, McMurray JG, Carey E, Hayes P, Feldman RA.

Human-to-human transmission of *Pseudomonas pseudomallei*.

Ann Intern Med. 1975 ;83:512-513.

[59] Mollaret HH.

« L'affaire du jardin des plantes » ou comment la mélioïdose fit son apparition en France.

Med Mal infect. 1988 (spécial Novembre):643-654.

[60] Olive C, Loetitia G, Desbois N, Roche B, Jouannelle J, Dodin A.

Forme septico-pyohémique de mélioïdose humaine : un premier cas aux Antilles Françaises.

Presse Med. 1995 ;24:1270.

[61] Osteraas GR, Hardman JM, Bass JW, Wilson C.

Neonatal melioidosis.

Am J Dis Child. 1971 ;122:446-448.

[62] Parry CM, Wuthiekanun V, Hoa NT, Diep TS, Thao LT, Loc PV, et al.

Melioidosis in Southern Vietnam : clinical surveillance and environmental sampling.

Clin Infect Dis. 1999 ;29:1323-1326.

[63] Peetermans WE, Van Wijngaerden E, Van Eldere J, Verhaegen J. Melioidosis brain and lung abscess after travel to Sri Lanka.

Clin Infect Dis. 1999 ;28:921-922.

[64] Perez JM, Petiot A, Adjide C, Gerry F, Goursaud R, Juminer B.

First case report of melioidosis in Guadeloupe, a French West Indies archipelago.

Clin Infect Dis. 1997 ;25:164-165.

[65] Perret JL.

La mélioïdose : une « bombe à retardement tropicale » en voie de dissémination ?

Med Trop. 1997 ;57:195-201.

[66] Perret JL, Vidal L, Thibault F.

La mélioïdose pulmonaire.

Rev Pneumol Clin. 1998 ;54:365-372.

- [67] Phetsouvanh R, Phongmany S, Newton P, Mayxay M, Ramsay A, Wuthiekanun V, et al.
Melioidosis and Pandora's box in the Lao People's Democratic Republic.
Clin Infect Dis. 2001 ;32(4):653-4.
- [68] Piggott JA, Hochholzer L.
Human melioidosis. A histopathologic study of acute and chronic melioidosis.
Arch Pathol. 1970 ;90:101-111.
- [69] Pongsunk S, Thirawattanasuk N, Piyasangthong N, Ekpo P.
Rapid identification of *Burkholderia pseudomallei* in blood cultures by a monoclonal antibody assay.
J Clin Microbiol. 1999 ;37:3662-3667.
- [70] Pons R, Advier M.
Melioidosis in Cochin China.
J Hyg. 1927 ;26:28-30.
- [71] Puthucheary SD, Parasakthi N, Lee MK.
Septicaemic melioidosis : a review of 50 cases from Malaysia.
Trans R Soc Trop Med Hyg. 1992 ;86:683-685.
- [72] Puthucheary SD, Vadivelu J, Ce-Cile, C, Kum-Thong, W, Ismail G.
Short report : Electron microscopic demonstration of extracellular structure of *Burkholderia pseudomallei*.
Am J Trop Hyg. 1996 ;54:313-314.
- [73] Ramsay SC, Labrooy J, Norton R, Webb B.
Demonstration of different patterns of musculoskeletal, soft tissue and visceral involvement in melioidosis using ^{99m}Tc stannous colloid white cell scanning.
Nucl Med Commun. 2001 ;22:1193-1199.
- [74] Sandford JP.
Pseudomonas species (including melioidosis and glanders) in Mandel GL, Bennet JE, Dolin R.
Principles and practice of infectious diseases. 4ème éd.- vol. 2-New-York : Churchill Livingstone,1995:2003-2006.

[75] Schwarzmaier A, Riezinger-Geppert F, Schober G, Karnik R, Valentin A. Fulminant septic melioidosis after a vacation in Thailand.

Wien Klin Wochenschr. 2000 ;112(20):892-5.

[76] Sexton MM, Jones AL, Chaowagul W, Woods DE.

Purification and characterization of a protease from *Pseudomonas pseudomallei*.

Can j Microbiol. 1994 ;40:903-910.

[77] Singcharoen T.

CT findings in melioidosis.

Australas Radiol. 1989 ;33:376-378.

[78] Sirisinha S, Anuntagool N, Dharakul T, Ekpo P, Wongratanacheewin S, Naigowit P et al.

Recent developments in laboratory diagnosis of melioidosis.

Acta Trop. 2000 ;74:235-245.

[79] Smith MD, Angus BJ, Wuthiekanun V, White NJ.

Arabinose assimilation defines a nonvirulent biotype of *Burkholderia pseudomallei*.

Infect Immun. 1997 ;65:4319-4321.

[80] Suputtamongkol Y, Chaowagul W, Chetchotisakd P, Lertpatanasuwun N, Intaranongpai S, Ruchutrakool T et al.

Risk factors for melioidosis and bacteremic melioidosis.

Clin Infect Dis. 1999 ;29:408-413.

[81] Suputtamongkol Y, Hall AJ, Dance DA, Chaowagul W, Rajchanuvong A, Smith MD et al.

The epidemiology of melioidosis in Ubon Ratchatani, northeast Thailand.

Int J Epidemiol. 1994 ;23:1082-1090.

[82] Suputtamongkol Y, Rajchanuwong A, Chaowagul W, Dance DA, Smith MD, Wuthiekanun V et al.

Ceftazidime vs. amoxicillin/clavulanate in the treatment of severe melioidosis. Clin Infect Dis. 1994 ;19:846-853.

[83] Stanton AT, Fletcher W.

Melioidosis.

- Studies from the Institute of Medical Research, federal Malay States. London : John Bale and Danielson,1932.
- [84] Steinmetz I, Rohde M, Brenneke B.
Purification and characterization of an exopolysaccharide of Burkholderia (Pseudomonas) pseudomallei.
Infect Immun. 1995 ;63(10):3959-65.
- [85] Tan AP, Pui MH, Tan LK.
Imaging patterns in melioidosis.
Australas Radiol. 1995 ;39:260-264.
- [86] Tan JK, Yip SK, Png DJ, Moorthy P.
Primary melioidotic prostatic abscess : presentation, diagnosis and management.
ANZ J Surg. 2002 ;72:408-410.
- [87] Thomas AD, Forbes-Faulkner JC.
Persistence of Pseudomonas pseudomallei in soil.
Aust Vet J. 1981 ;57:535-536.
- [88] Tiangpitayakorn C, Songsivilai S, Piyasangthong N, Dharakul T.
Speed of detection of Burkholderia pseudomallei in blood cultures and its correlation with the clinical outcome.
Am J Trop Med Hyg. 1997 ;57:96-99.
- [89] Turner MO, Lee VT, FitzGerald JM.
Melioidosis in a diabetic sailor.
Chest. 1994 ;106:952-954.
- [90] Van Phung L, Quynh HT, Yabuuchi E, Dance DA.
Pilot study of exposure to Pseudomonas pseudomallei in northern Vietnam.
Trans R Soc Trop Med Hyg. 1993 ;87:416.
- [91] Vasu C, Vadivelu J, Puthuchery SD.
The humoral immune response in melioidosis patients during therapy. Infection. 2003 ;31:24-30.
- [92] Veron M.

Pseudomonadaceae in. Le Minor L, Veron M eds.

Bactériologie médicale. Paris :Flammarion,1989:555-598.

[93] Visca P, Cazzola G, Petrucca A, Braggion C.

Travel-associated Burkholderia pseudomallei infection (Melioidosis) in a patient with cystic fibrosis : a case report.

Clin Infect Dis. 2001 Jan ;32(1) :E15-6. Epub 2000 Dec 11.

[94] Walsh AL, Smith MD, Wuthiekanun V, Suputtamongkol Y, Chaowagul W, Dance DA et al.

Prognostic significance of quantitative bacteremia in septicemic melioidosis. Clin Infect Dis. 1995 ;21:1498-1500.

[95] Walsh AL, Wuthiekanun V.

The laboratory diagnosis of melioidosis.

Br J Biomed Sci. 1996 ;53:249-253.

[96] White NJ.

Melioidosis in. Hunter GW, Strickland GT eds.

Hunter's tropical medicine and emerging infectious diseases. Philadelphia : Saunders WB,2000:313-316.

[97] White NJ.

Melioidosis.

Lancet. 2003 ;361:1715-1722.

[98] White NJ.

Melioidosis.

Zbl Bakt. 1994 ;280:439-443.

[99] Whitmore A, Krishnaswami CS.

An account of the discovery of a hitherto undescribed infective disease occurring among the population of Rangoon.

The Indian Medical Gazette. 1912 ;47:262-267.

[100] Wibulpolprasert B, Dhiensiri T.

Visceral organ abscesses in melioidosis : sonographic findings.

J Clin Ultrasound. 1999 ;27:29-34.

[101] Winstanley C, Hart CA.

Presence of type III secretion genes in *Burkholderia pseudomallei* correlates with Ara(-) phenotypes.

J Clin Microbiol. 2000 Feb ;38:883-885.

[102] Wong KT, Puthuchery SD, Vadivelu J.

The histopathology of human melioidosis.

Histopathology. 1995 ;26:51-55.

[103] Woods DE, DeShazer D, Moore RA, Brett PJ, Burtnick MN, Reckseidler SL et al.

Current studies on the pathogenesis of melioidosis.

Microbes Infect. 1999 ;1:157-162.

[104] Woods ML 2nd, Currie BJ, Howard DM, Tierney A, Watson A, Anstey NM et al.

Neurological melioidosis : seven cases from the Northern Territory of Australia.

Clin Infect Dis. 1992 ;15:163-169.

[105] Wuthiekanun V, Suputtamongkol Y, Simpson AJ, Kanaphun P, White NJ.

Value of throat swab in diagnosis of melioidosis.

J Clin Microbiol. 2001 ;39:3801-3802.

[106] Yabuuchi E, Kosako Y, Oyaizu H, Yano I, Hotta H, Hashimoto Y, et al.

Proposal of *Burkholderia* gen. nov. and transfer of seven species of the genus *Pseudomonas* homology group II to the new genus, with the type species *Burkholderia cepacia* (Palleroni and Holmes 1981) comb. nov.

Microbiol Immunol. 1992 ;36:1251-1275.

[107] Yang S.

Melioidosis research in China.

Acta Trop. 2000:157-165.

[108] Yip SK, Ang BS, Tan J.

Clinics in diagnostic imaging (57). Melioidotic prostatic abscess.

Singapore Med J. 2001 ;42:041-043.